

Assessment of Whole Brain White Matter Integrity in Youths and Young Adults With a Family History of Substance-Use Disorders

Ashley Acheson,^{1,2*} S. Andrea Wijtenburg,³ Laura M. Rowland,^{3,4}
Anderson M. Winkler,^{5,6} Frank Gaston,³ Charles W. Mathias,¹ Peter T. Fox,²
William R. Lovallo,⁷ Susan N. Wright,³ L. Elliot Hong,³
Donald M. Dougherty,¹ and Peter Kochunov³

¹Department of Psychiatry, University of Texas Health Science Center at San Antonio, San Antonio, Texas

²Research Imaging Institute, University of Texas Health Science Center at San Antonio, San Antonio, Texas

³Maryland Psychiatric Research Center, Department of Psychiatry, University of Maryland School of Medicine, Baltimore, Maryland

⁴Russell H. Morgan Department of Radiology and Radiological Science, Johns Hopkins University, Baltimore, Maryland

⁵Oxford Centre for Functional MRI of the Brain, University of Oxford, Oxford, United Kingdom

⁶Department of Psychiatry, Yale University School of Medicine, New Haven, Connecticut

⁷Behavioral Sciences Laboratories, Veterans Affairs Medical Center and University of Oklahoma Health Sciences Center, Oklahoma City, Oklahoma



Abstract: Individuals with a family history of substance use disorders (FH+) are at a greater risk of developing substance use disorders than their peers with no such family histories (FH−) and this vulnerability is proportional to the number of affected relatives (FH density). The risk for developing substance use disorders peaks during adolescence to early adulthood in the general population, and that is thought to be related to delayed maturation of frontocortical and frontostriatal functional circuits. We hypothesized that FH+ youth and young adults have impaired myelination of frontocortical and frontostriatal white matter tracts. We examined fractional anisotropy (FA) data in 80 FH+ and 34 FH− youths (12.9 ± 1.0 years) and in 25 FH+ and 30 FH− young adults (24.3 ± 3.4 years). FH+ youths had lower FA values in both frontocortical and frontostriatal tracts as well as parietocortical tracts including the anterior, superior and posterior corona radiata and the superior frontal-occipital fasciculus. Moreover, FA values in these tracts were negatively correlated with FH density. FH+ adults had lower

Additional Supporting Information may be found in the online version of this article.

Contract grant sponsor: NIDA, NIAAA, NIBIB, and NIMH of the National Institutes of Health; Contract grant numbers: R01-AA012207; R01-AA019691; R01-DA026868; R01-DA033997; R01-EB015611; R01-MH094520; T32-MH067533; Contract grant sponsor: Department of Veterans Affairs Medical Research Service; Contract grant number: CX000252

*Correspondence to: Ashley Acheson, Department of Psychiatry and Research Imaging Institute, The University of Texas Health

Science Center at San Antonio, 7703 Floyd Curl Drive, MC 6240, San Antonio, TX 78229. E-mail: acheson@uthscsa.edu

Received for publication 14 March 2014; Revised 8 May 2014; Accepted 13 May 2014.

DOI: 10.1002/hbm.22559

Published online 00 Month 2014 in Wiley Online Library (wileyonlinelibrary.com).

FA values in two frontocortical tracts: the genu of the corpus callosum and anterior corona radiata and also significant negative correlations between FA and FH density in these same tracts. In both groups, lower FA values corresponded to higher radial diffusivity suggesting reduced axonal myelination. We interpreted our findings as evidence for impaired myelination of frontal white matter that was proportional to FH density. Our data suggest that deficits may partially resolve with age, paralleling an age-related decline in risk for developing substance use disorders. *Hum Brain Mapp* 00:000–000, 2014. © 2014 Wiley Periodicals, Inc.

Key words: frontal white matter; family history; risk; diffusion tensor imaging; substance use

INTRODUCTION

Youths and young adults with a family history of alcohol and/or other drug use disorders (FH+) are at a greater risk for developing substance use disorders compared to their peers with no such history (FH−) [Finn et al., 1990; Lieb et al., 2002; McCaul et al., 1990; Merikangas et al., 1998]. Moreover, their risk of developing substance use disorders is proportional to the number of biological parents and grandparents with substance use disorders (FH density) [Dawson et al., 1992; Stoltenberg et al., 1998]. This increased risk has a genetic basis as demonstrated by twin, adoption, cross-fostering, and pedigree analysis studies [Cloninger et al., 1981; Merikangas 1990; Reich et al., 1998; Slutske et al., 2002], and is associated with a phenotypic pattern of “behavioral undercontrol” characterized by increased sensation seeking, risk-taking, aggressiveness, and antisocial behaviors [Sher et al., 2004; Sher and Trull, 1994; Tarter et al., 2003]. Likewise, FH+ individuals show deficits in executive functioning, impulse control, decision-making, and attention [Acheson et al., 2011a,b; Corral et al., 2003; Deckel 1999; Lovallo et al., 2006; Stevens et al., 2003] and altered activity in frontocortical and frontostriatal regions [Acheson et al., 2009; Cservenka et al., 2012; Glahn et al., 2007; Heitzeg et al., 2010; Schweinsburg et al., 2004; Silveri et al., 2011]. These findings suggest FH+ individuals have impairments in frontocortical and frontostriatal functioning, although specific mechanisms underlying this impairment and as well as potential relationships to risk for substance use disorders remain elusive.

The risk of developing a substance use disorder in FH+ individuals peaks during adolescence and early adulthood, similar to trends observed in general population [McGorry et al., 2011]. It has been hypothesized that this increased vulnerability in adolescence and early adulthood in the general population is driven at least in part by the relatively slow maturation of frontocortical and frontostriatal tracts [Blakemore and Robbins, 2012; Ernst and Fudge, 2009; Somerville and Casey, 2010], which not reach peak myelination levels until an individual is in their 30s or 40s, compared to the motor and sensory tracts that reach peak maturity in their teens or early 20s [Kochunov et al., 2011b, 2012; Sowell et al., 2003; Westlye et al., 2010]. This delayed maturation of

frontocortical and frontostriatal tracts is thought to lead to poor inhibition of reward-seeking and risk-taking behavior, including propensity toward problem substance use [Blakemore and Robbins, 2012; Ernst and Fudge, 2009; Somerville and Casey, 2010]. Plausibly, individuals more prone toward engaging in problem substance use in adolescence and early adulthood may have even more delayed or impaired frontocortical and frontostriatal white matter development during this period.

It is therefore plausible that FH+ individuals have delayed or impaired development of frontocortical and frontostriatal white matter relative to their peers and this contributes to the propensity toward problem substance use and other excessive reward seeking and risk taking behaviors. Consistent with this hypothesis, FH+ adolescents appear to show reduced myelination of frontocortical and frontostriatal pathways compared to FH− adolescents (Herting et al., 2010). However, this study had a small sample size (13 FH+ and 14 FH−) and was not sufficiently powered to examine additional relationships within the FH+ group, such as the impact of FH density. In addition, it is not clear whether FH+ related deficits in forebrain white matter persist past adolescence or are resolved during later development. To address these questions, we examined integrity of cerebral white matter in two large and well-characterized cohorts of FH+ youths and adults. These cohorts were collected as part of two independent study protocols and procedural differences prevented us from combining the cohorts. However, examining the two cohorts allows us to make initial observations about the possible course of FH+ related white matter deficits from preadolescence to early adulthood.

We estimated myelination levels using diffusion tensor imaging (DTI) to index fractional anisotropy (FA) of water diffusion. FA describes the directional selectivity of the random diffusion of water molecules [Basser 1994; Conturo et al., 1996; Pierpaoli and Basser, 1996; Ulug et al., 1995]. Higher FA values (maximum theoretical value is 1.0) are observed along heavily myelinated white matter tracts. The structure of the axonal cell membranes and myelin sheath hinders the diffusion of water molecules in all directions except along the fiber tract, therefore producing highly anisotropic water diffusion [Pierpaoli and Basser, 1996]. The absolute white matter FA values are sensitive to a variety of

parameters including regional myelination levels, the degree of intra-voxel fiber crossing, axonal density and average axonal diameter [Beaulieu, 2002]. However, changes in regional FA values during normal maturation and aging are shown to be predominantly due to changes in axonal myelination and can therefore be used as indirect measurement of myelin level [Budde et al., 2007; Madler et al., 2008; Song et al., 2003, 2005]. Several recent studies used FA of cerebral white matter to study heterocronicity of cerebral myelination and showed that the FA values of frontocortical and frontostriatal white matter tracts increase rapidly during the adolescence and suggest their peak myelination level is not reached until the an individual is in their twenties or thirties [Hasan et al., 2009a,b; Kochunov, et al., 2011b, 2012; McLaughlin et al., 2007].

In the present study, we tested two hypotheses. First, we hypothesized that FH+ individuals have impaired myelination of frontocortical and frontostriatal axonal tracts, as evidenced by the lower FA values. Second, we hypothesized that the degree of deficit in white matter integrity would be proportional to the FH density. We compared FA values in both cohorts across 14 major white tracts and specifically in frontocortical tracts such as the anterior corona radiata and frontostriatal tracts such as the superior frontooccipital fasciculus.

METHODS

Overview

We examined whole brain DTI data from two independent cohorts of FH+ and FH− individuals. The first cohort consisted of preadolescent to early adolescent youths and the second cohort consisted of young adults. Imaging data for both cohorts were collected using the same scanner and RF coil with similar high angular resolution DTI protocols and analyzed using identical procedures. Our statistical approach consisted of first comparing FH group differences across whole-brain average FA and regional values. Next, we examined correlations between FA and FH density. All analyses were repeated after controlling for possible demographic confounding variables. Finally, we also examined regional differences in the axial and radial diffusivities in secondary analyses. Reduced FA values can be caused by increased intravoxel crossing and tortuosity or twisting of white matter fibers rather than simply decreased myelination [Beaulieu, 2002]. However, parallel results with radial but not axial diffusivity would suggest FA differences are caused by differences in diffusivity across the axonal membranes, specifically implicating altered axonal myelination.

Youth Cohort

Participants

Thirty-four FH− and 80 FH+ youths (10 to 14 years old, average age = 12.9 ± 1.0) were recruited from a cohort of

386 volunteers in a community based ongoing longitudinal study of adolescent development and substance use involvement in youths at elevated risk for substance use disorders [Ryan et al., under review]. FH+ youths were deliberately oversampled to ensure a range in substance outcomes in the longitudinal portion of the study. Estimated intelligence was assessed using the Wechsler Abbreviated Scale of Intelligence [WASI; Psychological Corporation, 1999]. Family socioeconomic status was measured using the Four Factor Index of Socioeconomic Status [FFISS; Hollingshead, 1975]. Psychiatric symptoms and diagnoses was assessed using the Schedule for Affective Disorders and Schizophrenia for School-Age Children-Present and Lifetime Version [K-SADS-PL; Kaufman, et al., 1997] administered by trained research assistants and reviewed by a staff psychiatrist board-certified in child and adolescent psychiatry. Externalizing and internalizing symptoms for all participants were assessed from parent or guardian reports on the Child Behavior Checklist [Achenbach, 2001]. Exclusion criteria included: regular substance use [defined as substance use at least once per month for 6 consecutive months; Clark et al., 2005], positive breath alcohol or urine drug test at time of screening, low IQ (<70), implantable metallic devices or braces, or physical or developmental disabilities that would interfere with the subject's ability to understand or complete study requirements. Oppositional defiant disorder, conduct disorder, attention deficit/hyperactivity disorder, dysthymia or anxiety disorders were not exclusionary for FH+ in the youth cohort because these disorders are commonly comorbid with problem substance use [Iacono et al., 2008]. All participants signed consent forms approved by the Institutional Review Board of The University of Texas Health Science Center at San Antonio approved the study procedures. Privacy was further protected by a Certificate of Confidentiality from the Department of Health and Human Services.

Family history of substance use disorders

Family history classification for the youth cohort was established using the family history assessment module [Rice et al., 1995] based on parent report. All FH+ participants had a biological father with a past or present substance use disorder. Most FH+ youths (79%) had a father with an alcohol use disorder history, and 64% had a father with history of alcohol and other drug use disorders. Some FH+ youths (31%) had a biological mother with substance use disorder histories. Some (18%) had a mother with an alcohol use disorder history and 11% had a mother with history of alcohol and other drug use disorders. The most common other drug use disorders among parents were cannabis and stimulant use disorders. A family history density score (FH density) was calculated by counting the number of biological parents and grandparents meeting criteria for any substance use disorder. Affected grandparents had similar alcohol and other drug




Figure 1.

Significant correlation coefficients ($P \leq 0.05$, corrected) between regional FA values and FH density (number of parents and grandparents with substance use disorders) are color-coded for individual white matter tracts is shown for the youth cohort (left) and young adult cohort (right). Lateral (top) and medial (bottom) projections are shown.

use disorder histories as affected parents. Scores ranged from 0 (FH- participants) to a possible 6 (FH+ participants with both parents and all grandparents affected).

Collection and processing of magnetic resonance imaging data

All MRI procedures were performed at the Research Imaging Institute, University of Texas Health Science Center at San Antonio using Siemens Tim Trio 3T MR system (Erlangen, Germany) equipped with a multichannel head coil using a protocol described elsewhere [Wijtenburg et al., 2012]. A single-shot, echo-planar, single refocusing spin-echo sequence was used to acquire diffusion-weighted data with a spatial resolution of $1.7 \times 1.7 \times 3.0 \text{ mm}^3$. The sequence parameters were: TE/TR = 83/7,000 ms, FOV = 200 mm, two diffusion weighing values $b = 0$ and 700 s mm^{-2} and five $b = 0$ (non-diffusion weighted) images, 64 isotropically distributed diffusion-weighted directions, and axial slice orientation with 50

slices and no gaps. The number of directions, $b = 0$ images, and the magnitude of the b values were calculated using an optimization technique that maximizes the contrast to noise ratio based on the average diffusivity of the cerebral white matter and the T_2 relaxation times [Jones et al., 1999]. In addition, high-resolution T1-weighted data were collected using an optimized protocol described previously [Kochunov et al., 2006].

Image processing

DTI data was processed using a tract-based spatial statistics method, distributed as a part of FMRIB Software Library (FSL) package [Smith et al., 2006] as described elsewhere [Kochunov et al., 2011b]. The population-based, 3D, DTI cerebral white matter tract atlas developed at Johns Hopkins University and distributed with the FSL package [Wakana et al., 2004] was used to calculate population average FA values along the spatial course of 14, major white matter tracts (Table II, Fig. 1) as described

elsewhere [Kochunov et al., 2011a, 2012]. Briefly, The Johns Hopkins University atlas was nonlinearly aligned to the minimal deformation target brain and image containing labels for individual tracts was transferred to the minimal deformation target brain space using nearest-neighbor interpolation. Per-tract average values were calculated by averaging the values along the tracts in both hemispheres. The overall average FA values were calculated by averaging values for the entire white matter skeleton.

Adult Cohort

Participants

Thirty FH– and 25 FH+ young adults (18 to 30 years old, average age = 24.3 ± 3.4 years) were recruited from a larger cohort of 450 volunteers participating in a community based ongoing study examining behavioral and physiological characteristics of healthy young adults at elevated risk of alcohol and other substance use disorders [Lovallo et al., 2013]. Estimated intelligence (Shipley mental age) was determined from the vocabulary score on the Revised Shipley Institute of Living Scale [Zachary, 1986]. SES was measured using the Hollingshead scale [Hollingshead, 1975] with updated occupational categories and was based on the primary occupation of the main breadwinner in the household in which the participant grew up. Psychological functioning was assessed using the computerized version of the Diagnostic Interview Schedule-IV [DIS-IV; Bucholz et al., 1994] and the Beck Depression Inventory II [BDI; Beck et al., 1996]. Antisocial traits were quantified with the Sociability scale of the California Personality Inventory [CPI-So; Gough, 1994], a 46-item self-report measure of norm abiding and pro-social behaviors. Alcohol and drug use were assessed through the Cahalan Drinking Habits Questionnaire [Cahalan and Cisin, 1968], the alcohol use disorders identification test [Barbor et al., 1992], and a drug use questionnaire [Saunders et al., 2008]. Exclusion criteria included: a history of alcohol or drug dependence, substance abuse or depression within the past 2 months, a history of any other DSM-IV-TR (APA, 2000) Axis I disorder, failing a urine drug screen or breath-alcohol test on days of testing, or physical or developmental disabilities that would interfere with the subject's ability to understand or complete study requirements. All participants signed consent forms approved by the Institutional Review Boards of the University of Oklahoma Health Sciences Center and the Veterans Affairs Medical Center in Oklahoma City, OK and at the University of Texas Health Sciences Center, San Antonio, TX and were paid for their participation. Privacy was further protected by a Certificate of Confidentiality from the Department of Health and Human Services.

Family history of substance use disorders

Family history classification for the adult cohort was established using the family history research diagnostic

criteria [Andreasen et al., 1977; Zimmerman et al., 1988] and confirmed by parent report for all subjects. Almost all FH+ adults (80%) had a father with substance use disorder histories. Most (72%) had a father with an alcohol use disorder history, and 28% had a father with history of alcohol and other drug use disorders. Some FH+ adults (42%) had a biological mother with substance use disorder histories. Some (31%) had a mother with an alcohol use disorder history and 23% had a mother with history of alcohol and other drug use disorders. Information on the type of other drug use disorders was not recorded. Affected grandparents had similar alcohol and other drug use disorder histories as affected parents. FH density was calculated as in the youth cohort, with scores ranging from 0 to 6.

Diffusion tensor imaging and data processing

All imaging was also performed on the same scanner and the same sequence controls parameters. The only difference was that 55 isotropically distributed diffusion-weighted directions were used instead of 64, in the interest of shorter scan time. The adult cohort DTI data were processed identically to the youth cohort DTI data.

Statistics

FA values averaged for the whole brain and regional white matter measurements were compared between FH+ and FH– individuals in both cohorts using independent sample *t* tests. In addition, we examined linear correlations between regional FA values and FH density in both groups separately. We then replicated these same analyses examining radial and axial diffusivity in place of FA. If the FH differences are driven by differences in myelination, we would expect parallel results with radial but not axial diffusivity.

Binary logistic regressions were used to test for significant relationships between FH group and FA tracts while controlling for socioeconomic status and the presence of externalizing and internalizing disorders. In these analyses FH group was the dependent variable, and the predictors were identified as those tracts that significantly differentiated groups in the preceding *t*-test comparisons after Bonferroni correction. The regressions were conducted using individual models rather than multivariate because FA in all these tracts was significantly correlated with one another. A similar approach was used with linear regressions to predict FH density.

RESULTS

Youth Cohort

Participant characteristics

Demographic data for the youth cohort are summarized in Table I. The FH+ and FH– groups did not differ in

TABLE I. Demographic information for the youth cohort

	FH- (<i>n</i> = 34)		FH+ (<i>n</i> = 80)	
	Mean	(SD)	Mean	(SD)
Age	12.9	(1.1)	12.9	(1.2)
Wechsler abbreviated scale of intelligence	102.8	(12.2)	96.3	(12.5) ^a
Four factor index of socioeconomic status	43.6	(10.2)	34.0	(12.6) ^a
CBCL externalizing problems	2.0	(2.6)	7.4	(7.6) ^a
CBCL internalizing problems	2.9	(4.0)	6.4	(6.2) ^a
	Number	(%)	Number	(%)
Gender				
Male	19	(56)	39	(49)
Female	15	(44)	41	(51)
Race				
African-American	1	(3)	11	(14)
Caucasian	32	(94)	67	(84)
Other	1	(3)	2	(3)
Ethnicity				
Hispanic/Latino	29	(85)	63	(79)
Non-Hispanic/Latino	5	(15)	17	(21)
Externalizing disorders				
Attention deficit hyperactivity disorder	0	(0)	16	(20)
Conduct disorder	0	(0)	3	(4)
Oppositional defiant disorder	0	(0)	10	(13)
Internalizing disorders				
Generalized anxiety disorder	0	(0)	7	(9)
Separation anxiety disorder	0	(0)	2	(3)
Specific phobia	0	(0)	7	(9)
Lifetime alcohol and drug use (n. ever used)				
Alcohol	1	(3)	6	(8)
Marijuana	0	(0)	4	(5)
Tobacco	1	(3)	2	(3)
Other	0	(0)	0	(0)

FH+ = family history of substance use disorders (SUDs); FH- = no family history of SUDs; CBCL= Child Behavior Checklist (48).

^a*P* < 0.05.

age, race, or ethnicity or drug use. However, FH+ participants had lower IQ, socioeconomic status, and had more externalizing and internalizing problems.

Imaging findings

FA data were discarded from 5 FH- females, 3 FH+ females, and 2 FH+ males, due to movement artifacts (a spatial shift of over 3 mm between consecutive imaging frames). Post-hoc analyses demonstrated no significant differences in the average motion between FH+ and FH- individuals (average motion per TR = 0.41 ± 0.11 mm vs. 0.42 ± 0.12 mm, *P* = 0.95, for FH+ and FH- subjects, respectively).

FH+ youths had significantly lower FA values in fronto-cortical and frontostriatal tracts including the anterior and superior corona radiata and superior frontal-occipital fasciculus (all *P* < 0.05 with Bonferroni correction, Table II). Similarly, FH density was significantly and negatively correlated with FA values primarily in the same tracts,

indicating that progressively lower FA values were seen in individuals with more parents and grandparents with substance use disorders (Table II, Fig. 1, Supporting Information Fig. 1S). Similar deficits were observed in parietocortical tracts including the posterior corona radiata. These FA differences remained significant after excluding youths with externalizing and internalizing disorders (Supporting Information Table IS). Post-hoc analyses indicate that reduced FA in FH+ individuals due to elevations in the radial diffusivity (Supporting Information Table IIS). FH+ subjects also showed some modest reductions in axial diffusivity but most were not significant after controlling for multiple comparisons (Table IIIS, see Supporting Information).

FA values of the anterior and superior corona radiata were independent predictors of FH group after controlling for socioeconomic status and the presence of externalizing and internalizing disorders (Table III). Anterior corona radiata FA correctly classified 80% of the sample (57% of FH- and 89% of FH+), and FA in the superior corona

TABLE II. Youth cohort fractional anisotropy (FA) values

Tract	Mean (\pm SD)		Significance of group-wise <i>t</i> test <i>P</i>	FH density correlation	
	FH- (<i>n</i> = 30)	FH+ (<i>n</i> = 75)		<i>r</i>	<i>P</i>
Corpus Callosum					
Genu	0.763 (0.035)	0.746 (0.027)	0.009	-0.196	0.044
Body	0.664 (0.033)	0.656 (0.040)	0.314	-0.145	0.139
Splenium	0.771 (0.022)	0.763 (0.024)	0.121	-0.167	0.087
Internal capsule	0.635 (0.017)	0.625 (0.019)	0.013	-0.182	0.063
Corona radiata					
Anterior	0.479 (0.019)	0.463 (0.018)	1.8×10^{-5a}	-0.404	1.8×10^{-5a}
Superior	0.517 (0.015)	0.504 (0.019)	0.001 ^a	-0.332	5.1×10^{-4a}
Posterior	0.528 (0.017)	0.515 (0.021)	0.001 ^a	-0.307	8.7×10^{-4a}
Posterior thalamic radiation	0.637 (0.026)	0.623 (0.027)	0.020	-0.320	0.001 ^a
Sagittal stratum	0.574 (0.031)	0.563 (0.026)	0.076	-0.175	0.074
External capsule	0.494 (0.017)	0.489 (0.017)	0.202	-0.070	0.479
Cingulum	0.623 (0.028)	0.611 (0.028)	0.043	-0.152	0.122
Superior longitudinal fasciculus	0.514 (0.019)	0.505 (0.022)	0.043	-0.135	0.170
Frontal-occipital fasciculus					
Superior	0.543 (0.0278)	0.523 (0.027)	0.002 ^a	-0.342	3.6×10^{-4a}
Inferior	0.523 (0.038)	0.522 (0.040)	0.872	0.012	0.905

Uncorrected *P* values reported.

^a*P* ≤ 0.05 with Bonferroni correction (*P* ≤ 0.004).

radiata correctly classified 79% of the sample (57% of FH- and 88% of FH+). Similarly, FA in the anterior, superior, and posterior corona radiata, thalamic radiation, and superior frontal-occipital fasciculus all were significant predictors of density of family history of substance use disorders, even after controlling for socioeconomic status and the presence of externalizing and internalizing disorders.

Adult Cohort

Participant characteristics

The FH+ and FH- groups did not differ in age, estimated intelligence, race, or ethnicity, but FH+ participants had a significantly lower socioeconomic status (Table IV).

Imaging findings

No data from adults had to be excluded due to excessive movement and there were no difference in the average motion between FH+ and FH- individuals (average motion per TR = 0.39 ± 0.13 mm vs. 0.38 ± 0.12 mm, *P* = 0.95, for FH+ and FH- subjects, respectively). FH+ adults had lower FA values only in frontocortical tracts, specifically the genu of the corpus callosum and anterior corona radiata, and FH density was significantly negatively correlated with FA values in these same tracts (Table V, Fig. 1, Supporting Information Fig. 2S). Similar to the youth cohort, radial diffusivity was elevated in the frontocortical tracts in the FH+ cohort (Table IVS). There were no significant group differences in the axial diffusivity (Table VS). In addition, we observed a significant

TABLE III. Regression analyses for youth cohort

Tract	Family history binary logistic regression			Nagelkerke <i>R</i> ²	Family density linear regression			
	Wald	<i>P</i>	OR 95% CI		Standardized β	<i>t</i>	<i>P</i>	<i>R</i> ²
Corona radiata								
Anterior	10.12	0.001	0.995 (0.993–0.998)	0.394	-0.350	-4.10	<0.001	0.281
Superior	9.63	0.002	0.995 (0.992–0.998)	0.395	-0.345	-4.01	<0.001	0.280
Posterior	-	-	-	-	-0.268	-3.04	0.003	0.232
Thalamic radiation	-	-	-	-	-0.334	-3.93	<0.001	0.273
Superior frontal-occipital fasciculus	-	-	-	-	-0.347	-4.12	<0.001	0.282

TABLE IV. Demographic information in the adult cohort

	FH- (<i>n</i> = 30)		FH+ (<i>n</i> = 25)	
	Mean	(SD)	Mean	(SD)
Age	24.5	(2.7)	24.1	(4.2)
Shipley vocabulary score	31	(4.3)	30	(4.2)
Hollingshead and Redlich socioeconomic status score	49	(13)	40	(15) ^a
California personality inventory sociability scale score	32.1	(4.0)	30.2	(4.3)
Beck depression inventory score	4.0	(5.7)	4.7	(5.3)
Age at first drink	17	(2.9)	15	(4.0)
Drinks per month	35.4	(46.6)	19.8	(29.1)
	Number	(%)	Number	(%)
Gender				
Male	13	(43)	7	(28)
Female	17	(57)	18	(72)
Race ^a				
African-American	0	(0)	7	(28)
Caucasian	27	(90)	18	(72)
Other	3	(10)	0	(0)
Ethnicity				
Hispanic/Latino	3	(10)	4	(16)
Non-Hispanic/Latino	27	(90)	21	(84)
Tobacco smoker	1	(3)	1	(4)
Past alcohol abuse	3	(11)	1	(4)
Past other substance use disorders	0	(0)	0	(0)
Lifetime illicit drug use (n ever used)				
Marijuana	13	(4)	14	(5)
Other drug use	6	(20)	8	(32)

^a*P* < 0.05.

TABLE V. Adult cohort fractional anisotropy (FA) values

Tract	Mean (\pm SD)		Significance of group-wise <i>t</i> test <i>P</i>	FH density correlation	
	FH- (<i>n</i> = 30)	FH+ (<i>n</i> = 25)		<i>r</i>	<i>P</i>
Corpus callosum					
Genu	0.767 (0.029)	0.739 (0.054)	0.017	-0.399	0.003 ^a
Body	0.685 (0.040)	0.679 (0.048)	0.586	-0.125	0.365
Splenium	0.780 (0.023)	0.779 (0.029)	0.850	-0.005	0.969
Internal capsule	0.626 (0.023)	0.629 (0.027)	0.673	0.101	0.461
Corona radiata					
Anterior	0.539 (0.023)	0.518 (0.033)	0.010	-0.397	0.003 ^a
Superior	0.510 (0.022)	0.507 (0.026)	0.729	-0.110	0.425
Posterior	0.517 (0.023)	0.517 (0.021)	0.964	-0.029	0.832
Posterior thalamic radiation	0.620 (0.027)	0.620 (0.015)	0.955	-0.018	0.895
Sagittal stratum	0.574 (0.029)	0.573 (0.027)	0.869	0.098	0.477
External capsule	0.498 (0.021)	0.496 (0.024)	0.807	-0.041	0.764
Cingulum	0.637 (0.035)	0.633 (0.031)	0.652	-0.133	0.334
Superior longitudinal fasciculus	0.511 (0.025)	0.511 (0.019)	0.938	-0.132	0.338
Frontal-occipital fasciculus					
Superior	0.573 (0.042)	0.572 (0.043)	0.923	-0.120	0.382
Inferior	0.492 (0.045)	0.495 (0.056)	0.865	0.030	0.827

Uncorrected *P* values reported.

^a*P* ≤ 0.05 with Bonferroni correction (*P* ≤ 0.004).

correlation between FH density and thalamic radiation axial diffusivity.

Anterior corona radiata FA was a significant predictor of FH density, even after controlling for socioeconomic status (standardized $\beta = -0.363$; $t = -2.89$; $P = 0.006$, $R^2 = 0.200$).

DISCUSSION

Our study demonstrated significantly reduced fractional anisotropy (FA) of cerebral white matter in both the youth and young adult FH+ individuals. FH+ youth had lower FA in frontocortical and frontostriatal tracts, and the degree of impairment was proportional to the number of affected parents and grandparents, an empirical risk factor for developing substance use disorder (Dawson et al., 1992; Stoltenberg et al., 1998). FH+ youth also had reduced FA in parietocortical tracts, and FA was negatively correlated with FH density in these same tracts. In FH+ adults, FA was decreased only in frontocortical tracts, and FA was negatively correlated to FH density in these tracts as well. Analyses of radial and axial diffusivities indicate the reduced FA values in FH+ individuals were caused by higher radial diffusivity (across the axonal membranes), specifically implicating reduced axonal myelination. Our findings in adolescent cohort provide strong support for the hypothesis that adolescent FH+ individuals may experience altered or delayed development of frontal white matter. Our findings in young adults suggests that white matter deficits may partially be resolved by early adulthood, paralleling a decline in the risk for developing substance use disorders from adolescence to adulthood.

Our findings replicate and extend the previous study that identified FA deficits in a small sample of FH+ youths [Herting et al., 2010]. We replicated the findings of lower FA values in frontal white matter of FH+ in both youth and adult cohorts. We also replicated the earlier study's finding of lower parietocortical FA in FH+ youth but did not observe this in FH+ adults. Moreover, we demonstrated that FA deficits were proportional to FH density in both youths and young adults. Similarly, others have reported reduced frontal as well as parietal FA values in individuals with substance use disorders [Chung et al., 2007; Lim et al., 2008; Lin et al., 2012; Liu et al., 2010; Ma et al., 2009; McQueeney et al., 2009; Romero et al., 2010; Upadhyay et al., 2010]. Collectively, these findings suggest that impaired integrity of the frontal white matter may be a neurobiological phenotype associated with having a family history of substance use disorders. Furthermore, these findings demonstrate that white matter deficits in FH+ individuals are present before and may potentially contribute to the development of substance use disorders. Longitudinal studies will be needed to directly evaluate this hypothesis, and we are prospectively following youth cohort to investigate these issues. In fact, our cohort was chosen to oversample the FH+ youths and

include those with externalizing and internalizing conditions to increase the likelihood of observing a range of substance use outcomes.

Further, our study demonstrated the reduced FA values in FH+ individuals generally corresponded with increased radial but not axial diffusivity, suggesting that reduced axonal myelination is the likely biological mechanism behind our FA findings. Methodologically, the absolute FA values are only indirect measurements of axonal myelination as FA is also sensitive to factors such as the axonal diameter, intra-voxel changes in fiber orientation and others [Budde et al., 2007; Song et al., 2003, 2005]. Radial diffusivity is a measure of a restricted diffusivity across the axonal walls. Biologically, this measure indexes the permeability of axonal membranes and therefore serves as an indirect estimate of axonal myelination level. Furthermore, studies in animal models where levels of myelination were carefully manipulated showed that changes in the radial diffusivity (e.g., diffusivity across the axonal membranes) was responsible for changes in FA values [Budde et al., 2007; Gao et al., 2009; Song et al., 2003, 2005]. Consistent with this rational, radial diffusivity values for the tracts that showed significantly lower FA in FH+ group were also significantly and positively correlated with the FH density. This was in contrast with the axial diffusivity measurements for the same areas that showed sporadic, weaker relationships with FH group status and FH density.

The regional pattern of group-wise FA differences are consistent with the heterochronicity of cerebral myelination trends, first reported by Fleschsig [1901], however the biological basis of our findings cannot be fully understood because the biology of the regional heterochronicity during normal cerebral myelination is not well known. The pattern of regional differences between FH+ youth and their FH- peers pointed specifically at the frontocortical and frontostriatal tracts as well as parietocortical tracts that carry multimodal, associative and projection fibers with no group differences for the FA values of the sensory and motor tracts. Microscopy studies show that oligodendrocytes that myelinate associative tracts are morphological distinct from these that myelinate motor and sensory tracts [Pfefferbaum et al., 2000; Sulliva et al., 2001; Wood and Bunge, 1984]. The former type of oligodendrocytes produce far fewer myelin layers per axon (less myelin) and have reduced rates of myelin turn over and slower rates of myelin repair than the glia located in the primary sensory and motor tracts [Hof et al., 1990; Lamantia and Rakic, 1990; Wakana et al., 2004]. Recent studies indicate there is strong genetic control over normal cerebral myelination process [Kanchibhotla et al., 2013]. Therefore, a plausible explanation for our findings is the genotype-by-age interactions that affect cerebral development that may be regionally specific. Recently advanced gene-by-environment interaction analysis demonstrated that genes influence neurocognitive traits and white matter coherence as a function of age from young adulthood to old age

[Glahn et al., 2013]. In another example, polymorphisms in the *TP53*, a gene central to DNA repair were specifically associated with deficits in the multimodal, frontal matter integrity WM [Molina et al., 2011]. The specific genes that exert risk for developing alcohol and other substance use disorders are unknown but this risk has a strong genetic basis [Cloninger et al., 1981; Merikangas, 1990; Reich et al., 1998; Slutske et al., 2002]. Therefore, our findings may help future genetic and genotype-by-age studies to choose quantitative endophenotypes to get measurements closely linked with the neurobiology of this disorder.

The white matter deficits in FH+ adolescents were more pervasive and included frontocortical, frontostriatal, and parietocortical tracts compared to FH+ adults where FA deficits were confined to frontocortical tracts. This difference is unlikely to be brought about by the procedural differences as imaging data for both cohorts were collected on the same scanner and RF coil using very similar protocols and analyzed using identical procedures. Similarly, the youth cohort findings were not the result of including individuals externalizing and internalizing disorders since the same findings were observed after excluding these individuals. In fact, this may suggest that the decreased frontocortical, frontostriatal, and parietocortical white matter in FH+ individuals may be caused by delays in the normal maturation of these tracts during development. This also implies that many FH+ individuals eventually reach normal myelination levels at adulthood, at least in some tracts. A relatively limited window of impaired white matter development in adolescence through early adulthood in FH+ individuals would parallel the enhanced risk for developing substance use disorders during this period [McGorry et al., 2011], and may plausibly contribute to in the initiation and progression of substance use disorders. However, larger cross sectional studies, including a broader age range of participants, and prospective studies following FH+ youth into adulthood are needed to test this hypothesis.

Our study has both strengths and limitations. Strengths include the use of two independent, well-characterized, large sample study cohorts tested with nearly identical DTI protocols. It is unlikely that the findings were a consequence of substance use as the adult cohort was free of substance use disorders and none of the youth cohort had any history of regular substance use or and most had not even having tried alcohol or other drugs. Limitations include a lack of prospective data on substance use outcomes, although the youth cohort is currently being followed longitudinally. Although the results suggest some white matter impairments observed in FH+ youths may be resolved by adulthood, we were not able to directly compare findings across the cohorts and specifically test for age effects due to the slight differences in the DTI protocol. Additionally our youth cohort was predominantly Hispanic, while our adult cohort was predominantly non-Hispanic with relatively fewer males. Finally, it possible that by not testing FH+ adults with substance use

disorders we may have restricted the adult cohort to individuals less predisposed to substance use disorders, although non-affected FH+ individuals may still have inherited risk factors [Cloninger et al., 1981], similar to other complex genetic disorders where individuals may have inherited predisposing factors without ultimately becoming afflicted themselves [Falconer, 1965; Falconer, 1967]. Thus it possible that factors other than development may have contributed to the different findings between the two cohorts. Finally, reduced FA values can be caused by increased intravoxel crossing and tortuosity or twisting of white matter fibers [Beaulieu, 2002]. However, in this study, we consider altered myelination to be the likely culprit for reduced FA values because of the parallel findings with radial, but not axial, diffusivity.

CONCLUSIONS

In the present study, we observed impairments in frontocortical, frontostriatal and parietocortical white matter in FH+ youths that were robustly associated with FH density, before initiation of any regular drug or alcohol use, and obtained similar FA findings limited to frontocortical white matter in a young adult cohort. Our findings suggest that FH+ youths have impaired development of frontal and parietal white matter that may at least partially resolve by early adulthood, potentially paralleling their increased risk for substance use disorder development in adolescence through early adulthood. However, additional cross-sectional and longitudinal studies will be needed to directly investigate this possibility as well as examine relationships with frontal white matter deficits and the initiation and progression of substance use disorders.

ACKNOWLEDGMENTS

The content is solely the view of the authors and does not necessarily represent the official view of the National Institutes of Health or the VA. Marika Vela-Gude, David Hernandez, Nora Charles, Anran Xu, and Andrew Cohoon provided excellent technical assistance.

REFERENCES

- Achenbach TM (2001): Child behavior checklist for ages 6 to 18. Burlington, VT: University of Vermont, Research Center for Children, Youth, and Families.
- Acheson A, Robinson JL, Glahn DC, Lovallo WR, Fox PT (2009): Differential activation of the anterior cingulate cortex and caudate nucleus during a gambling simulation in persons with a family history of alcoholism: Studies from the Oklahoma Family Health Patterns Project. *Drug Alcohol Depend* 100:17–23.
- Acheson A, Richard DM, Mathias CW, Dougherty DM (2011a): Adults with a family history of alcohol related problems are more impulsive on measures of response initiation and response inhibition. *Drug Alcohol Depend* 117:198–203.

- Acheson A, Vincent AS, Sorocco KH, Lovallo WR (2011b): Greater discounting of delayed rewards in young adults with family histories of alcohol and drug use disorders: Studies from the Oklahoma family health patterns project. *Alcohol Clin Exp Res* 35:1607–1613.
- Andreasen NC, Endicott J, Stitzer RL, Winokur G (1977): The family history method using diagnostic criteria. *Arch Gen Psychiatry* 34:1229–1235.
- APA (2000): *Diagnostic and Statistical Manual of Mental Disorders: DSM-IV-TR*. Washington, DC: American Psychiatric Association.
- Barbor TF, de las Fuentes JR, Saunders JB, Grant M (1992): AUDIT: The Alcohol Use Disorders Identification Test: Guidelines for use in Primary Health Care. Geneva, Switzerland: World Health Organization.
- Basser PJ (1994): Focal magnetic stimulation of an axon. *IEEE Trans Biomed Eng* 41:601–606.
- Beaulieu C (2002): The basis of anisotropic water diffusion in the nervous system—A technical review. *NMR Biomed* 15:435–455.
- Beck AT, Steer RA, Ball R, Ranieri W (1996): Comparison of beck depression inventories -IA and -II in psychiatric outpatients. *J Pers Assess* 67:588–597.
- Blakemore SJ, Robbins TW (2012): Decision-making in the adolescent brain. *Nat Neurosci* 15:1184–1191.
- Bucholz KK, Cadoret R, Cloninger CR, Dinwiddie SH, Hesselbrock VM, Nurnberger JI Jr, Reich T, Schmidt I, Schuckit MA (1994): A new, semi-structured psychiatric interview for use in genetic linkage studies: A report on the reliability of the SSAGA. *J Stud Alcohol* 55:149–158.
- Budde MD, Kim JH, Liang HF, Schmidt RE, Russell JH, Cross AH, Song SK (2007): Toward accurate diagnosis of white matter pathology using diffusion tensor imaging. *Magn Reson Med* 57:688–695.
- Cahalan D, Cisin IH (1968): American drinking practices: Summary of findings from a national probability sample: II. Measurement of massed versus spaced drinking. *Quart J Stud Alcohol* 29:642–656.
- Chung A, Lyoo IK, Kim SJ, Hwang J, Bae SC, Sung YH, Sim ME, Song IC, Kim J, Chang KH, Renshaw PF (2007): Decreased frontal white-matter integrity in abstinent methamphetamine abusers. *Int J Neuropsychopharmacol* 10:765–775.
- Clark DB, Cornelius JR, Kirisci L, Tarter RE (2005): Childhood risk categories for adolescent substance involvement: A general liability typology. *Drug Alcohol Depend* 77:13–21.
- Cloninger CR, Bohman M, Sigvardsson S (1981): Inheritance of alcohol abuse: Cross fostering analysis of adopted men. *Arch Gen Psychiatry* 38:861–868.
- Conturo TE, McKinstry RC, Akbudak E, Robinson BH (1996): Encoding of anisotropic diffusion with tetrahedral gradients: A general mathematical diffusion formalism and experimental results. *Magn Reson Med* 35:399–412.
- Corporation P (1999): *Psychological Corporation, Wechsler Abbreviated Scale of Intelligence (WASI) Manual*. San Antonio, TX: Harcourt Brace and Company.
- Corral M, Holguin SR, Cadaveira F (2003): Neuropsychological characteristics of young children from high-density alcoholism families: A three-year follow-up. *J Stud Alcohol* 64:195–199.
- Cservenka A, Herting MM, Nagel BJ (2012): Atypical frontal lobe activity during verbal working memory in youth with a family history of alcoholism. *Drug Alcohol Depend* 123:98–104.
- Dawson DA, Harford TC, Grant BF (1992): Family history as a predictor of alcohol dependence. *Alcohol Clin Exp Res* 16:572–575.
- Deckel AW (1999): Tests of executive functioning predict scores on the MacAndrew Alcoholism Scale. *Prog Neuropsychopharmacol Biol Psychiatry* 23:209–223.
- Ernst M, Fudge JL (2009): A developmental neurobiological model of motivated behavior: Anatomy, connectivity and ontogeny of the triadic nodes. *Neurosci Biobehav Rev* 33:367–382.
- Falconer DS (1965): The inheritance of liability to certain diseases, estimated from the incidence among relatives. *Ann Hum Genet Lond* 29:51–76.
- Falconer DS (1967): The inheritance of liability to diseases with variable age of onset, with particular reference to diabetes mellitus. *Ann Hum Genet* 31:1–20.
- Finn PR, Kleinman I, Pihl RO (1990): The lifetime prevalence of psychopathology in men with multigenerational family histories of alcoholism. *J Nervous Mental Dis* 178:500–504.
- Flechsigs P (1901): Developmental (myelogenetic) localisation of the cerebral cortex in the human. *Lancet* 158:1027–1030.
- Gao W, Lin W, Chen Y, Gerig G, Smith JK, Jewells V, Gilmore JH (2009): Temporal and spatial development of axonal maturation and myelination of white matter in the developing brain. *AJNR Am J Neuroradiol* 30:290–296.
- Glahn DC, Lovallo WR, Fox PT (2007): Reduced amygdala activation in young adults at high risk of alcoholism: Studies from the Oklahoma family health patterns project. *Biol Psychiatry* 61:1306–1309.
- Glahn DC, Kent JW Jr, Sprooten E, Diego VP, Winkler AM, Curran JE, McKay DR, Knowles EE, Carless MA, Goring HH, Dyer TD, Olvera RL, Fox PT, Almasy L, Charlesworth J, Kochunov P, Duggirala R, Blangero J (2013): Genetic basis of neurocognitive decline and reduced white-matter integrity in normal human brain aging. *Proc Natl Acad Sci USA* 110:19006–19011.
- Gough HG (1994): Theory, development, and interpretation of the CPI socialization scale. *Psychol Rep* 75:651–700.
- Hasan KM, Iftikhar A, Kamali A, Kramer LA, Ashtari M, Cirino PT, Papanicolaou AC, Fletcher JM, Ewing-Cobbs L (2009a): Development and aging of the healthy human brain uncinate fasciculus across the lifespan using diffusion tensor tractography. *Brain Res* 1276:67–76.
- Hasan KM, Kamali A, Iftikhar A, Kramer LA, Papanicolaou AC, Fletcher JM, Ewing-Cobbs L (2009b): Diffusion tensor tractography quantification of the human corpus callosum fiber pathways across the lifespan. *Brain Res* 1249:91–100.
- Heitzeg MM, Nigg JT, Yau WY, Zucker RA, Zubieta JK (2010): Striatal dysfunction marks preexisting risk and medial prefrontal dysfunction is related to problem drinking in children of alcoholics. *Biol Psychiatry* 68:287–295.
- Herting MM, Schwartz D, Mitchell SH, Nagel BJ (2010): Delay discounting behavior and white matter microstructure abnormalities in youth with a family history of alcoholism. *Alcohol Clin Exp Res* 34:1590–1602.
- Hof PR, Cox K, Morrison JH (1990): Quantitative analysis of a vulnerable subset of pyramidal neurons in Alzheimer's disease: I. Superior frontal and inferior temporal cortex. *J Comp Neurol* 301:44–54.
- Hollingshead AB (1975): *Four Factor Index of Social Status*. New Haven, CT: Yale University. p 22.

- Iacono WG, Malone SM, McGue M (2008): Behavioral disinhibition and the development of early-onset addiction: Common and specific influences. *Annu Rev Clin Psychol* 4:325–348.
- Jones DK, Horsfield MA, Simmons A (1999): Optimal strategies for measuring diffusion in anisotropic systems by magnetic resonance imaging. *Magn Reson Med* 42:515–525.
- Kanchibhotla SC, Mather KA, Wen W, Schofield PR, Kwok JB, Sachdev PS (2013): Genetics of ageing-related changes in brain white matter integrity—A review. *Ageing Res Rev* 12:391–401.
- Kaufman J, Birmaher B, Brent D, Rao U, Flynn C, Moreci P, Williamson D, Ryan N (1997): Schedule for affective disorders and schizophrenia for school-age children-present and lifetime version (K-SADS-PL): Initial reliability and validity data. *J Am Acad Child Adolesc Psychiatry* 36:980–988.
- Kochunov P, Lancaster JL, Glahn DC, Purdy D, Laird AR, Gao F, Fox P (2006): Retrospective motion correction protocol for high-resolution anatomical MRI. *Hum Brain Mapp* 27:957–962.
- Kochunov P, Glahn D, Nichols T, Winkler A, Hong E, Holcomb H, Stein J, Thompson P, Curran J, Carless M, Olvera R, Johnson M, Cole S, Kochunov V, Kent J, Blangero J (2011a): Genetic analysis of cortical thickness and fractional anisotropy of water diffusion in the brain. *Front Neurosci* 5:1–15.
- Kochunov P, Glahn DC, Lancaster J, Thompson PM, Kochunov V, Rogers B, Fox P, Blangero J, Williamson DE (2011b): Fractional anisotropy of cerebral white matter and thickness of cortical gray matter across the lifespan. *Neuroimage* 58:41–49.
- Kochunov P, Williamson DE, Lancaster J, Fox P, Cornell J, Blangero J, Glahn DC (2012): Fractional anisotropy of water diffusion in cerebral white matter across the lifespan. *Neurobiol Aging* 33:9–20.
- Lamantia AS, Rakic P (1990): Cytological and quantitative characteristics of four cerebral commissures in the rhesus monkey. *J Comp Neurol* 291:520–537.
- Lieb R, Merikangas KR, Hofler M, Pfister H, Isensee B, Wittchen HU (2002): Parental alcohol use disorders and alcohol use and disorders in offspring: A community study. *Psychol Med* 32: 63–78.
- Lim KO, Wozniak JR, Mueller BA, Franc DT, Specker SM, Rodriguez CP, Silverman AB, Rotrosen JP (2008): Brain macrostructural and microstructural abnormalities in cocaine dependence. *Drug Alcohol Depend* 92:164–172.
- Lin WC, Chou KH, Chen CC, Huang CC, Chen HL, Lu CH, Li SH, Wang YL, Cheng YF, Lin CP (2012): White matter abnormalities correlating with memory and depression in heroin users under methadone maintenance treatment. *PloS One* 7: e33809.
- Liu IC, Chiu CH, Chen CJ, Kuo LW, Lo YC, Tseng WY (2010): The microstructural integrity of the corpus callosum and associated impulsivity in alcohol dependence: A tractography-based segmentation study using diffusion spectrum imaging. *Psychiatry Res* 184:128–134.
- Lovallo WR, Yechiam E, Sorocco KH, Vincent AS, Collins FL (2006): Working memory and decision-making biases in young adults with a family history of alcoholism: Studies from the Oklahoma family health patterns project. *Alcoholism Clin Exp Res* 30:763–773.
- Lovallo WR, Farag NH, Sorocco KH, Acheson A, Cohoon AJ, Vincent AS (2013): Early life adversity contributes to impaired cognition and impulsive behavior: Studies from the Oklahoma family health patterns project. *Alcohol Clin Exp Res* 37:616–623.
- Ma L, Hasan KM, Steinberg JL, Narayana PA, Lane SD, Zuniga EA, Kramer LA, Moeller FG (2009): Diffusion tensor imaging in cocaine dependence: Regional effects of cocaine on corpus callosum and effect of cocaine administration route. *Drug Alcohol Depend* 104:262–267.
- Madler B, Drabycz SA, Kolind SH, Whittall KP, MacKay AL (2008): Is diffusion anisotropy an accurate monitor of myelination? Correlation of multicomponent T2 relaxation and diffusion tensor anisotropy in human brain. *Magn Reson Imaging* 26:874–888.
- McCaul ME, Turkkan JS, Svikis DS, Bigelow GE, Cromwell CC (1990): Alcohol and drug use by college males as a function of family alcoholism history. *Alcohol Clin Exp Res* 14:467–471.
- McGorry PD, Purcell R, Goldstone S, Amminger GP (2011): Age of onset and timing of treatment for mental and substance use disorders: Implications for preventive intervention strategies and models of care. *Curr Opin Psychiatry* 24:301–306.
- McLaughlin NC, Paul RH, Grieve SM, Williams LM, Laidlaw D, DiCarlo M, Clark CR, Whelihan W, Cohen RA, Whitford TJ, Gordon E (2007): Diffusion tensor imaging of the corpus callosum: A cross-sectional study across the lifespan. *Int J Dev Neurosci* 25:215–221.
- McQueeny T, Schweinsburg BC, Schweinsburg AD, Jacobus J, Bava S, Frank LR, Tapert SF (2009): Altered white matter integrity in adolescent binge drinkers. *Alcohol Clin Exp Res* 33: 1278–1285.
- Merikangas KR (1990): The genetic epidemiology of alcoholism. *Psychol Med* 20:11–22.
- Merikangas KR, Stolar M, Stevens DE, Goulet J, Preisig MA, Fenton B, Zhang H, O'Malley SS, Rounsaville BJ (1998): Familial transmission of substance use disorders. *Arch Gen Psychiatry* 55:973–979.
- Molina V, Papiol S, Sanz J, Rosa A, Arias B, Fatjo-Vilas M, Calama J, Hernandez AI, Becker J, Fananas L (2011): Convergent evidence of the contribution of TP53 genetic variation (Pro72Arg) to metabolic activity and white matter volume in the frontal lobe in schizophrenia patients. *Neuroimage* 56:45–51.
- Pfefferbaum A, Sullivan EV, Hedehus M, Lim KO, Adalsteinsson E, Moseley M (2000): Age-related decline in brain white matter anisotropy measured with spatially corrected echo-planar diffusion tensor imaging. *Magn Reson Med* 44:259–268.
- Pierpaoli C, Basser PJ (1996): Toward a quantitative assessment of diffusion anisotropy. *Magn Reson Med* 36:893–906.
- Reich T, Edenberg HJ, Goate A, Williams JT, Rice JP, Van Eerdewegh P, Foroud T, Hesselbrock V, Schuckit MA, Bucholz K, Porjesz B, Li TK, Conneally PM, Nurnberger JI Jr, Tischfield JA, Crowe RR, Cloninger CR, Wu W, Shears S, Carr K, Crose C, Willig C, Begleiter H (1998): Genome-wide search for genes affecting the risk for alcohol dependence. *Am J Med Genet* 81: 207–215.
- Rice JP, Reich T, Bucholz KK, Neuman RJ, Fishman R, Rochberg N, Hesselbrock VM, Nurnberger JI Jr, Schuckit MA, Begleiter H (1995): Comparison of direct interview and family history diagnoses of alcohol dependence. *Alcohol Clin Exp Res* 19: 1018–1023.
- Romero MJ, Asensio S, Palau C, Sanchez A, Romero FJ (2010): Cocaine addiction: Diffusion tensor imaging study of the inferior frontal and anterior cingulate white matter. *Psychiatry Res* 181:57–63.
- Ryan SR, Acheson A, Charles NE, Lake SL, Hernandez DL, Mathias CW, Dougherty DM (Under Review): Clinical and

- social/environmental characteristics in a community sample of children with and without family histories of substance use disorder in the San Antonio area: A descriptive study.
- Saunders B, Farag N, Vincent AS, Collins FL Jr, Sorocco KH, Lovallo WR (2008): Impulsive errors on a Go-NoGo reaction time task: Disinhibitory traits in relation to a family history of alcoholism. *Alcohol Clin Exp Res* 32:888–894.
- Schweinsburg AD, Paulus MP, Barlett VC, Killeen LA, Caldwell LC, Pulido C, Brown SA, Tapert SF (2004): An FMRI study of response inhibition in youths with a family history of alcoholism. *Ann N Y Acad Sci* 1021:391–394.
- Sher KJ, Trull TJ (1994): Personality and disinhibitory psychopathology: Alcoholism and antisocial personality disorder. *J Abnormal Psychol* 103:92–102.
- Sher KJ, Grekin ER, Williams NA (2004): The development of alcohol use disorders. *Annu Rev Clin Psychol* 22:1–22.
- Silveri MM, Rogowska J, McCaffrey A, Yurgelun-Todd DA (2011): Adolescents at risk for alcohol abuse demonstrate altered frontal lobe activation during Stroop performance. *Alcohol Clin Exp Res* 35:218–228.
- Slutske WS, Heath AC, Madden PA, Bucholz KK, Statham DJ, Martin NG (2002): Personality and the genetic risk for alcohol dependence. *J Abnorm Psychol* 111:124–133.
- Smith SM, Jenkinson M, Johansen-Berg H, Rueckert D, Nichols TE, Mackay CE, Watkins KE, Ciccarelli O, Cader MZ, Matthews PM, Behrens TE (2006): Tract-based spatial statistics: Voxelwise analysis of multi-subject diffusion data. *Neuroimage* 31:1487–1505.
- Somerville LH, Casey BJ (2010): Developmental neurobiology of cognitive control and motivational systems. *Curr Opin Neurobiol* 20:236–241.
- Song SK, Sun SW, Ju WK, Lin SJ, Cross AH, Neufeld AH (2003): Diffusion tensor imaging detects and differentiates axon and myelin degeneration in mouse optic nerve after retinal ischemia. *Neuroimage* 20:1714–1722.
- Song SK, Yoshino J, Le TQ, Lin SJ, Sun SW, Cross AH, Armstrong RC (2005): Demyelination increases radial diffusivity in corpus callosum of mouse brain. *Neuroimage* 26:132–140.
- Sowell ER, Peterson BS, Thompson PM, Welcome SE, Henkenius AL, Toga AW (2003): Mapping cortical change across the human life span. *Nat Neurosci* 6:309–315.
- Stevens MC, Kaplan RF, Hesselbrock VM (2003): Executive-cognitive functioning in the development of antisocial personality disorder. *Addict Behav* 28:285–300.
- Stoltenberg SF, Mudd SA, Blow FC, Hill EM (1998): Evaluating measures of family history of alcoholism: Density versus dichotomy. *Addiction* 93:1511–1520.
- Sullivan EV, Adalsteinsson E, Hedehus M, Ju C, Moseley M, Lim KO, Pfefferbaum A (2001): Equivalent disruption of regional white matter microstructure in ageing healthy men and women. *Neuroreport* 12:99–104.
- Tarter RE, Kirisci L, Mezzich A, Cornelius JR, Pajer K, Vanyukov M, Gardner W, Blackson T, Clark D (2003): Neurobehavioral disinhibition in childhood predicts early age at onset of substance use disorder. *Am J Psychiatry* 160:1078–1085.
- Ulug AM, Barker PB, van Zijl PC (1995): Correction of motional artifacts in diffusion-weighted images using a reference phase map. *Magn Reson Med* 34:476–480.
- Upadhyay J, Maleki N, Potter J, Elman I, Rudrauf D, Knudsen J, Wallin D, Pendse G, McDonald L, Griffin M, Anderson J, Nutile L, Renshaw P, Weiss R, Becerra L, Borsook D (2010): Alterations in brain structure and functional connectivity in prescription opioid-dependent patients. *Brain* 133:2098–2114.
- Wakana S, Jiang H, Nagae-Poetscher LM, van Zijl PC, Mori S (2004): Fiber tract-based atlas of human white matter anatomy. *Radiology* 230:77–87.
- Westlye LT, Walhovd KB, Dale AM, Bjornerud A, Due-Tønnessen P, Engvig A, Grydeland H, Tamnes CK, Ostby Y, Fjell AM (2010): Differentiating maturational and aging-related changes of the cerebral cortex by use of thickness and signal intensity. *Neuroimage* 52:172–185.
- Wijtenburg SA, McGuire SA, Rowland LM, Sherman PM, Lancaster JL, Tate DF, Hardies LJ, Patel B, Glahn DC, Hong LE, Fox PT, Kochunov P (2012): Relationship between fractional anisotropy of cerebral white matter and metabolite concentrations measured using (1)H magnetic resonance spectroscopy in healthy adults. *Neuroimage* 66C:161–168.
- Wood P, Bunge RP (1984): The biology of the oligodendrocyte. In: Norton WT, editor. *Oligodendroglia*. New York: Plenum Press. pp 1–46.
- Zachary WT (1986): Shipley Institute of Living Scale Revised manual. Los Angeles: Western Psychological Services.
- Zimmerman M, Coryell W, Pfohl B, Stangl D (1988): The reliability of the family history method for psychiatric diagnoses. *Arch Gen Psychiatry* 45:320–322.